

UNCLASSIFIED

Defense Technical Information Center
Compilation Part Notice

ADP011973

TITLE: Constructive Approximation by $[V,f]$ -Reproducing Kernels

DISTRIBUTION: Approved for public release, distribution unlimited

This paper is part of the following report:

TITLE: International Conference on Curves and Surfaces [4th], Saint-Malo, France, 1-7 July 1999. Proceedings, Volume 2. Curve and Surface Fitting

To order the complete compilation report, use: ADA399401

The component part is provided here to allow users access to individually authored sections of proceedings, annals, symposia, etc. However, the component should be considered within the context of the overall compilation report and not as a stand-alone technical report.

The following component part numbers comprise the compilation report:

ADP011967 thru ADP012009

UNCLASSIFIED

Constructive Approximation by (V,f)-Reproducing Kernels

Mohammed-Najib Benbourhim

Abstract. In this paper we propose a constructive method to build reproducing kernels. We define the notion of (V, f) -reproducing kernel, and prove that every reproducing kernel is a (V, f) -reproducing kernel. We study the minimal approximation by these (V, f) -reproducing kernels for different choices of V and f . Examples to which our results apply include curve and surface fitting.

§1. (V, f) -Reproducing Kernels

For any set (respectively locally compact set) Ω , we denote by \mathbb{R}^Ω (respectively $C^m(\Omega)$) the space of real-valued functions (respectively m -times continuously differentiable functions) defined on Ω equipped with the topology of pointwise convergence (respectively uniform convergence on the compact subsets of Ω). Let us recall some definitions.

Definition 1.1. A real-valued function H defined on $\Omega \times \Omega$ is a reproducing kernel on $\Omega \times \Omega$ if

- 1) H is symmetric: $H(t, s) = H(s, t)$ for all $t, s \in \Omega$,
- 2) H is of positive type:

$$\sum_{k, l=1}^{k, l=N} \lambda_k \lambda_l H(t_k, t_l) \geq 0,$$

for any finite point set $\{t_k\}_{k=1}^N$ of Ω and real numbers $\{\lambda_k\}_{k=1}^N$.

Definition 1.2. A vector subspace \mathcal{H} of \mathbb{R}^Ω is said to be a hilbertian subspace of \mathbb{R}^Ω (respectively $C^m(\Omega)$) if

- 1) \mathcal{H} is a Hilbert space,
- 2) The natural injection from \mathcal{H} into \mathbb{R}^Ω (respectively $C^m(\Omega)$) is continuous.

We review some important results on reproducing kernels which are studied in [4].

Theorem 1.1.

- 1) A Hilbert space \mathcal{H} (respectively a real-valued function H defined on $\Omega \times \Omega$) is a hilbertian subspace of \mathbb{R}^Ω (respectively a reproducing kernel on $\Omega \times \Omega$) if and only if there exists one and only one reproducing kernel H on $\Omega \times \Omega$ (respectively hilbertian subspace \mathcal{H} of \mathbb{R}^Ω) such that

$$u(t) = \langle u \mid H(\cdot, t) \rangle_{\mathcal{H}}, \quad \forall t \in \Omega, \quad \forall u \in \mathcal{H}.$$

\mathcal{H} is called the hilbertian subspace associated with H .

- 2) For any hilbertian basis $(f_i)_{i \in I}$ of \mathcal{H} : $H(t, s) = \sum_{i \in I} f_i(t)f_i(s)$.
- 3) If H is separately m -times continuously differentiable, then \mathcal{H} is a hilbertian subspace of $\mathcal{C}^m(\Omega)$.
- 4) The vector space $\mathcal{H}_0 = \text{span}\{(H(\cdot, t))_{t \in \Omega}\}$ is dense in \mathcal{H} .

Let $(V, \langle \cdot \mid \cdot \rangle_V)$ be a Hilbert space, Ω be a set and f be a function from Ω into V .

Definition 1.3. For all $f : \Omega \longrightarrow V$, we define a (V, f) -reproducing kernel H_f by

$$H_f(t, s) = \langle f(t) \mid f(s) \rangle_V, \quad \forall (t, s) \in \Omega \times \Omega. \quad (1.1)$$

We have the following result:

Theorem 1.2. H_f defined by (1.1) is a reproducing kernel on $\Omega \times \Omega$ and its associated hilbertian subspace \mathcal{H}_f of \mathbb{R}^Ω is

$$\mathcal{H}_f = \left\{ w \in \mathbb{R}^\Omega \mid \exists u \in V : w(t) = \langle u \mid f(t) \rangle_V, \forall t \in \Omega \right\}.$$

Proof: One can easily verify that H_f is a reproducing kernel.

Let $\tilde{H}_f : V \longrightarrow \mathbb{R}^\Omega$ be defined by $(\tilde{H}_f u)(t) = \langle u \mid f(t) \rangle_V$. The mapping \tilde{H}_f is linear, and the inequality

$$|(\tilde{H}_f u)(t)| \leq \|u\|_V \|f(t)\|_V = \|u\|_V \|H_f(t, t)\|_V^{\frac{1}{2}},$$

for all $t \in \Omega$ and for all $u \in V$ implies that it is continuous. Let \mathcal{M} be the closure in V of the vector space $\text{span}\{(f(t))_{t \in \Omega}\}$, and $P_{\mathcal{M}}$ the orthogonal projector on \mathcal{M} . We define on $\mathcal{H}_f = \tilde{H}_f(V)$ the bilinear form

$$\langle \tilde{H}_f u \mid \tilde{H}_f v \rangle_{\mathcal{H}_f} = \langle P_{\mathcal{M}} u \mid P_{\mathcal{M}} v \rangle_V.$$

It is easy to see that this form is a scalar product on \mathcal{H}_f . Then the linear mapping $\tilde{H}_f : \mathcal{M} \longrightarrow \mathcal{H}_f$ is an isometry, and consequently $(\mathcal{H}_f, \langle \cdot \mid \cdot \rangle_{\mathcal{H}_f})$ is a Hilbert space. For all $t \in \Omega$, the function

$$H_f(t, \cdot) : s \in \Omega \longrightarrow H_f(t, s) = \langle f(t) \mid f(s) \rangle_V,$$

is an element of \mathcal{H}_f , and satisfies the reproducing formula

$$(\tilde{H}_f u)(t) = \langle \tilde{H}_f u \mid H_f(t, \cdot) \rangle_{\mathcal{H}_f}, \quad \forall u \in V.$$

Consequently (see Theorem 1.1), \mathcal{H}_f is a hilbertian subspace of \mathbb{R}^Ω and admits H_f as reproducing kernel. \square

Theorem 1.3. Let ω be a set and c a mapping from ω to Ω . Then

$$(Hc)_f(y, z) = H_f(c(y), c(z)) = \langle f(c(y)) \mid f(c(z)) \rangle_V$$

is a reproducing kernel on $\omega \times \omega$.

Proof: For all $y \in \omega$, $f(c(y))$ is in V . The function $(Hc)_f$ is symmetric and is of positive type:

$$\sum_{k,l=1}^{k=N} \lambda_k \lambda_l (Hc)_f(y_k, y_l) = \left\langle \sum_{k=1}^{k=N} \lambda_k f(c(y_k)) \mid \sum_{k=1}^{k=N} \lambda_k f(c(y_k)) \right\rangle_V \geq 0. \quad \square$$

Example 1.1. Let $V = L^2(a, b)$, Ω a subset of \mathbb{R} and $f(t)(x) = \exp(cxt)$ where c is a real constant. Then

$$H_f(t, s) = \begin{cases} (\exp(cb(t+s)) - \exp(ca(t+s)))/(c(t+s)), & \text{if } (t+s) \neq 0, \\ b-a, & \text{otherwise.} \end{cases}$$

Example 1.2. Let $V = L^2(\mathbb{R}^+)$, and suppose Ω is a subset of \mathbb{R}^n .

For all functions $c : \Omega \longrightarrow (0, +\infty)$, we have

$$(i) \text{ If } f(t)(x) = \frac{2}{\pi^{\frac{1}{2}}} \exp^{-c(t)|x|^2}, \text{ then } H_f(t, s) = \frac{1}{\sqrt{c(t) + c(s)}}.$$

$$(ii) \text{ If } f(t)(x) = \exp^{-c(t)x}, \text{ then } H_f(t, s) = \frac{1}{c(t) + c(s)}, \text{ and in particular if}$$

$c(t) = \frac{P(t)}{Q(t)}$ (with $P(t)$ and $Q(t)$ polynomials), we obtain the rational reproducing kernel

$$H_f(t, s) = \frac{Q(t)Q(s)}{P(t)Q(s) + P(s)Q(t)}.$$

§2. (V,f)-Reproducing Kernels of Convolution Type

We consider the case where

$$1) V = L^2(\mathbb{R}^n) \text{ and } \Omega = \mathbb{R}^n.$$

$$2) f(t)(x) = f(t-x), \text{ with } f \text{ in the familiar Sobolev space } H^m(\mathbb{R}^n).$$

$$\text{Then } H_f(t, s) = \int_{\mathbb{R}^n} f(t-x)f(s-x)dx.$$

Theorem 2.1. We have the following properties:

- 1) $H_f(t, s) = h_f(t - s)$ with $h_f(\xi) = (f * \check{f})(\xi) = \mathcal{F}(|\mathcal{F}f|^2)(\xi)$, where $\check{f}(x) = f(-x)$ and $\mathcal{F}f$ is the Fourier transform of f .
- 2) $h_f \in \mathcal{C}_0^m(\mathbb{R}^n) = \left\{ u \in \mathcal{C}^m(\mathbb{R}^n) \mid \lim_{|t| \rightarrow \infty} (D^\alpha u)(t) = 0, \quad 0 \leq |\alpha| \leq m \right\}$.
- 3) The associated hilbertian subspace of H_f is

$$\mathcal{H}_f = f * L^2(\mathbb{R}^n) \hookrightarrow \mathcal{C}_0^m(\mathbb{R}^n) \text{ (continuous embedding).}$$

- 4) In particular, if $|\mathcal{F}f| > 0$, then

$$\mathcal{H}_f = \left\{ w \in \mathcal{S}' \mid \mathcal{F}w \in L_{loc}^1(\mathbb{R}^n), \quad \frac{\mathcal{F}w}{\mathcal{F}f} \in L^2(\mathbb{R}^n) \right\},$$

equipped with the scalar product

$$\langle w_1 \mid w_2 \rangle_{\mathcal{H}_f} = \int_{\mathbb{R}^n} \frac{\mathcal{F}w_1(\xi) \overline{\mathcal{F}w_2(\xi)}}{|\mathcal{F}f(\xi)|^2} d\xi.$$

- 5) If f is radial, then h_f is radial: $H_f(t, s) = h_f(|t - s|)$.

- 6) For all distinct points $\{t_k\}_{k=1}^N$ in \mathbb{R}^n , the matrix $H_N = (H_f(t_k, t_l))_{1 \leq k, l \leq N}$ is invertible (strictly positive definite).

Proof:

- 1) We have

$$H_f(t, s) = (2\pi)^{-\frac{n}{2}} \int_{\mathbb{R}^n} e^{-i(t-s|\xi)} |\mathcal{F}f(\xi)|^2 d\xi = \mathcal{F}(|\mathcal{F}f|^2)(t - s).$$

- 2) $f \in H^m(\mathbb{R}^n) \Rightarrow D^\alpha h_f = (D^\alpha f) * \check{f} \in \mathcal{C}_0^0(\mathbb{R}^n)$ for $0 \leq |\alpha| \leq m$, (see [2]).

- 3) is a consequence of Theorem 1.2 and the property given in 1).

- 4) Since $\mathcal{H}_f \hookrightarrow \mathcal{C}_0^m(\mathbb{R}^n) \hookrightarrow \mathcal{S}'$, we have the equivalences:

$$\begin{aligned} \{w \in \mathcal{H}_f\} &\Leftrightarrow \{\exists u \in L^2(\mathbb{R}^n) : \mathcal{F}w = \mathcal{F}u\mathcal{F}f\} \\ &\Leftrightarrow \{\mathcal{F}w \in L_{loc}^1(\mathbb{R}^n), \quad \frac{\mathcal{F}w}{\mathcal{F}f} \in L^2(\mathbb{R}^n)\}. \end{aligned}$$

From Theorem 1.2, we have

$$w(t) = \langle u \mid f(t - \cdot) \rangle_{L^2(\mathbb{R}^n)} = \langle P_{\mathcal{M}}u \mid f(t - \cdot) \rangle_{L^2(\mathbb{R}^n)} = (P_{\mathcal{M}}u * f)(t).$$

Then $\mathcal{F}w = \mathcal{F}P_{\mathcal{M}}u\mathcal{F}f$, and

$$\begin{aligned} \langle w_1 \mid w_2 \rangle_{\mathcal{H}_f} &= \int_{\mathbb{R}^n} P_{\mathcal{M}}u_1(x) P_{\mathcal{M}}u_2(x) dx \\ &= \int_{\mathbb{R}^n} \mathcal{F}P_{\mathcal{M}}u_1(\xi) \overline{\mathcal{F}P_{\mathcal{M}}u_2(\xi)} d\xi = \int_{\mathbb{R}^n} \frac{\mathcal{F}w_1(\xi) \overline{\mathcal{F}w_2(\xi)}}{|\mathcal{F}f(\xi)|^2} d\xi. \end{aligned}$$

5) For any orthogonal matrix A ,

$$\begin{aligned} h_f(At) &= \int_{\mathbb{R}^n} f(x)f(At-x)dx = \int_{\mathbb{R}^n} f(Ax)f(A(t-x))dx \\ &= \int_{\mathbb{R}^n} f(x)f(t-x)dx = h_f(t), \end{aligned}$$

since $f(Ax) = f(x)$ and $|\det A| = 1$.

6) We suppose that $f \neq 0$ in $L^2(\mathbb{R}^n)$. Since the matrix

$H_N = \left(\int_{\mathbb{R}^n} f(t_k - x)f(t_l - x)dx \right)_{1 \leq k, l \leq N}$ is a Gram matrix, it is invertible if and only if the system $\{f(t_k - \cdot)\}_{k=1}^N$ is linearly independent in $L^2(\mathbb{R}^n)$. If $\sum_{k=1}^N c_k f(t_k - x) = 0$ in $L^2(\mathbb{R}^n)$, for $c_k \in \mathbb{R}$, $1 \leq k \leq N$, then by the Fourier transform we get $\left(\sum_{k=1}^N c_k e^{-i\langle t_k, \xi \rangle} \right) \mathcal{F}f(\xi) = 0$ in $L^2(\mathbb{R}^n)$.

The Lebesgue measure of the set $\mathcal{N} = \{\xi \in \mathbb{R}^n \mid \sum_{k=1}^N c_k e^{-i\langle t_k, \xi \rangle} = 0\}$ is equal to zero. Then $\mathcal{F}f$ vanishes outside \mathcal{N} , i.e: $\mathcal{F}f \equiv 0$ in $L^2(\mathbb{R}^n)$ and by the inverse Fourier transform, $f \equiv 0$ in $L^2(\mathbb{R}^n)$, which complete the proof. \square

Example 2.1. Let $u(x) = (1 - |x|)_+$ and $\mathcal{F}u = v$. We have $v(x) = \frac{\sin^2(\frac{x}{2})}{x^2}$.

(i) Taking $f = \mathcal{F}(|u|^{\frac{1}{2}})$, $H_f(t, s) = (1 - |t - s|)_+$.

(ii) Taking $f = \mathcal{F}(|v|^{\frac{1}{2}})$, $H_f(t, s) = \frac{\sin^2(\frac{t-s}{2})}{(t-s)^2}$.

Example 2.2. (Bessel reproducing kernels) For $n \in \mathbb{N}$, $\alpha \in \mathbb{R}$ and $\alpha > n$; consider $G_\alpha \in L^2(\mathbb{R}^n)$ defined by $\mathcal{F}(G_\alpha)(x) = (1 + |x|^2)^{-\frac{\alpha}{2}}$.

(i) Taking $f = \mathcal{F}(|G_\alpha|^{\frac{1}{2}})$, $H_f(t, s) = (1 + |t - s|^2)^{-\frac{\alpha}{2}}$.

(ii) Taking $f = \mathcal{F}(|\mathcal{F}G_{n+1}|^{\frac{1}{2}})$, $H_f(t, s) = \frac{\pi^{\frac{1-n}{2}}}{2^n \Gamma(\frac{n+1}{2})} \exp(-|t - s|)$ and

$\mathcal{H}_f = H^{\frac{n+1}{2}}(\mathbb{R}^n)$ (Sobolev space).

Example 2.3. (ν -B-spline reproducing kernels) Let

1) $Y_l(x) = \frac{1}{l!} x_+^l$.

2) $\nu \in \mathcal{E}'$ (distributions with compact support) such that $\nu(p) = 0$, for all polynomial p in $\mathcal{P}_l(\mathbb{R})$.

3) $f = \nu * Y_l$.

For such functions f , we give the following theorem without proof.

Theorem 2.2. We have:

- 1) $f \in L^2(\mathbb{R})$.
- 2) For all u in $V^{l+1}(\mathbb{R}) = \{v \in L^2_{loc}(\mathbb{R}) / v^{(l+1)} \in L^2(\mathbb{R})\}$ (Beppo-Levi space) we have $\int_{\mathbb{R}} u^{(l+1)}(x)f(t-x)dx = (\nu * u)(t)$.
- 3) $H_f(t, s) = (-1)^l(\check{\nu} * \nu * Y_{2l+1})(t-s)$ and $\mathcal{H}_f = \nu * V^{l+1}(\mathbb{R})$.

In the particular case of divided differences, ν is defined as the m th-iterated convolution $\nu = \frac{(\delta_a - \delta_b)^{*m}}{(b-a)}$, and $\check{\nu} = \frac{(\delta_{-a} - \delta_{-b})^{*m}}{(b-a)}$.

§3. Data Fitting by (V,f)-Reproducing Kernels

Let $\{t_k\}_{k=1}^N$ a set of distinct points in Ω , and define a linear operator A_N from \mathcal{H}_f into \mathbb{R}^N by $A_N(u) = (u(t_k))_{1 \leq k \leq N}$.

Definition 3.1. For all $z_N \in \mathbb{R}^N$ and $\epsilon \in [0, 1]$ we define a spline to be any solution of the following minimal approximation problem:

$$(P_\epsilon(z_N)) : \quad \inf_{u \in \mathcal{C}_\epsilon} \left((1 - \epsilon) \langle u | u \rangle_{\mathcal{H}_f} + \epsilon \|A_N u - z_N\|_{\mathbb{R}^N}^2 \right),$$

where

$$\mathcal{C}_\epsilon = \begin{cases} A_N^{-1}\{z_N\}, & \text{if } \epsilon = 0 \text{ (Interpolation),} \\ \mathcal{H}_f, & \text{if } \epsilon \in]0, 1[\text{ (Smoothing).} \end{cases}$$

The following theorem gives the spline in the case $\epsilon \neq 0$.

Theorem 3.1. For all $(\epsilon, z_N) \in]0, 1[\times \mathbb{R}^N$, the problem $P_\epsilon(z_N)$ (Smoothing) admits a unique solution

$$\sigma^\epsilon(t) = \sum_{k=1}^{k=N} \lambda_k^\epsilon H_f(t, t_k),$$

where the coefficients ${}^t\Lambda^\epsilon = (\lambda_1^\epsilon, \dots, \lambda_N^\epsilon) \in \mathbb{R}^N$ are the solution of the system

$$(H_N + \frac{\epsilon}{1-\epsilon} I_N) \Lambda_N^\epsilon = z_N,$$

with $H_N = (H_f(t_k, t_l))_{1 \leq k, l \leq N}$ and I_N is the identity matrix.

Proof: 1) From the continuous embedding: $\mathcal{H}_f \hookrightarrow \mathbb{R}^\Omega$ (see Theorem 1.2), we deduce that A_N is continuous. 2) $A_N(\mathcal{H}_f)$ is closed as a vector subspace of \mathbb{R}^N . Then from the general spline theory (see [1,3]) we get the theorem. \square

Theorem 3.2. *The following two properties are equivalent:*

- 1) For all $z_N \in \mathbb{R}^N$, the problem $P_0(z_N)$ admits a unique solution.
- 2) The system $\{f(t_k)\}_{k=1}^N$ is linearly independent in V .

Proof: For all $z_N \in \mathbb{R}^N$, the problem $P_0(z_N)$ admits a unique solution if and only if the matrix $H_N = (H_f(t_k, t_l))_{1 \leq k, l \leq N}$ is invertible (see [1,3]). Since the matrix $H_N = (\langle f(t_k) | f(t_l) \rangle_V)_{1 \leq k, l \leq N}$ is a Gram matrix, it is invertible if and only if the system $\{f(t_k)\}_{k=1}^N$ is linearly independent in V . \square

Furthermore, for the particular case $V = L^2(\mathbb{R}^n)$ and $f \in H^m(\mathbb{R}^n)$, Theorem 3.2 and the property (6) of Theorem 2.1 imply the following theorem.

Theorem 3.3. *For all $f \in H^m(\mathbb{R}^n)$ and $z_N \in \mathbb{R}^N$, the problem $P_0(z_N)$ (Interpolation) admits a unique solution*

$$\sigma^0(t) = \sum_{k=1}^{k=N} \lambda_k^0 H_f(t, t_k),$$

where the coefficients ${}^t\Lambda^0 = (\lambda_1^0, \dots, \lambda_N^0) \in \mathbb{R}^N$ are the solution of the system

$$H_N \Lambda_N^0 = z_N,$$

with $H_N = (H_f(t_k, t_l))_{1 \leq k, l \leq N}$ and I_N is the identity matrix.

§4. Data Fitting Preserving Polynomials

Let $\mathcal{P}_d(\mathbb{R}^n)$ the vector space of polynomials of degree at most d . We suppose:

(H1) For all $p \in \mathcal{P}_d(\mathbb{R}^n)$ the subset $\{t_k\}_{k=1}^N$ of \mathbb{R}^n is such that

$$\{p(t_k) = 0, \quad 1 \leq k \leq N\} \iff p \equiv 0.$$

(H2) $\mathcal{H}_f \cap \mathcal{P}_d(\mathbb{R}^n) = \{0\}$.

We remark that in the case $V = L^2(\mathbb{R}^n)$ and $f \in H^m(\mathbb{R}^n)$, the hypothesis (H2) is satisfied because $\mathcal{H}_f \subset C_0^m(\mathbb{R}^n)$ (see Theorem 2.1(2)), and

$$C_0^m(\mathbb{R}^n) \cap \mathcal{P}_d(\mathbb{R}^n) = \{0\}.$$

Let \mathcal{H}_f^d be the Hilbert direct sum: $\mathcal{H}_f^d = \mathcal{H}_f \oplus \mathcal{P}_d(\mathbb{R}^n)$. We denote by Π_f the orthogonal projector from \mathcal{H}_f^d onto \mathcal{H}_f , and we define on \mathcal{H}_f^d the linear mapping $A_N(u) = (u(t_k))_{1 \leq k \leq N} \in \mathbb{R}^N$. For all $(\epsilon, z_N) \in [0, 1] \times \mathbb{R}^N$, we consider the following minimal approximation problem in \mathcal{H}_f^d :

$$(P_\epsilon(z_N)) : \quad \inf_{u \in \mathcal{C}_\epsilon} \left((1 - \epsilon) \langle \Pi_f(u) | \Pi_f(u) \rangle_{\mathcal{H}_f} + \epsilon \|A_N u - z_N\|_{\mathbb{R}^N}^2 \right),$$

where

$$C_\epsilon = \begin{cases} A_N^{-1}\{z_N\}, & \text{if } \epsilon = 0 \text{ (Interpolation),} \\ \mathcal{H}_f^d, & \text{if } \epsilon \in]0, 1[\text{ (Smoothing),} \\ \mathcal{P}_d(\mathbb{R}^n), & \text{if } \epsilon = 1 \text{ (Least squares).} \end{cases}$$

The hypothesis (H1) implies that the problem $P_1(z_N)$ admits a unique solution. In the case $\epsilon \in]0, 1[$ (Smoothing) we have the following theorem:

Theorem 4.1. For all $(\epsilon, z_N) \in]0, 1[\times \mathbb{R}^N$, the problem $P_\epsilon(z_N)$ admits a unique solution σ^ϵ . In the case $\epsilon \in]0, 1[$, the solution σ^ϵ is given by

$$\sigma^\epsilon(t) = \sum_{k=1}^{k=N} \lambda_k^\epsilon H_f(t, t_k) + \sum_{i=1}^{i=n_d} b_i^\epsilon p_i(t),$$

where the coefficients ${}^t\Lambda^\epsilon = (\lambda_1^\epsilon, \dots, \lambda_N^\epsilon) \in \mathbb{R}^N$ and ${}^tB^\epsilon = (b_1^\epsilon, \dots, b_{n_d}^\epsilon) \in \mathbb{R}^{n_d}$ are the solution of the system

$$\begin{pmatrix} H_N + \frac{\epsilon-1}{\epsilon} I_N & E \\ {}^tE^\epsilon & 0 \end{pmatrix} \begin{pmatrix} \Lambda^\epsilon \\ B^\epsilon \end{pmatrix} = \begin{pmatrix} z_N \\ 0 \end{pmatrix},$$

with

1) $H_N = (H_f(t_k, t_l))_{1 \leq k, l \leq N}$ and I_N is the identity matrix,

2) $E = (E_{k,i})_{1 \leq i \leq n_d}^{1 \leq k \leq N}$ with $E_{k,i} = p_i(t_k)$ and $(p_i)_{1 \leq i \leq n_d}$ is a basis of $\mathcal{P}_d(\mathbb{R}^n)$.

In particular, if there exists $p \in \mathcal{P}_d(\mathbb{R}^n)$ such that $\{p(t_k) = z_{N,k}, 1 \leq k \leq N\}$, then $\sigma^\epsilon = p$ (preserving polynomials property).

Proof: Theorem 4.1 is a consequence of general spline theory (see [1,3]):

- 1) A_N is continuous since \mathcal{H}_f^d is a hilbertian subspace of \mathbb{R}^Ω .
- 2) Π_f is continuous and $\Pi_f(\mathcal{H}_f^d) = \mathcal{H}_f$ is closed since Π_f is an orthogonal projector.
- 3) $\ker A_N \cap \ker \Pi_f = \{0\}$: derives from the hypothesis (H1) and the fact that $\ker \Pi_f = \mathcal{P}_d(\mathbb{R}^n)$.
- 4) $\ker A_N + \ker \Pi_f$ is closed since $\ker \Pi_f$ is a finite dimensional vector space.

□

References

1. Attéa, M., *Hilbertian Kernels and Splines Functions*, Elsevier Science, North Holland, 1992.
2. Donoghue, W. F., *Distributions and Fourier Transforms*, Academic Press, 1969.
3. Laurent, P.-J., *Approximation et Optimisation*, Hermann, Paris, 1972.
4. Schwartz, L., Sous-espaces hilbertiens d'espaces vectoriels topologiques et noyaux associés, J. Analyse Math. **13** (1964), 115–256.

Benbourhim Mohammed-Najib
 Université Paul Sabatier (M.I.P)
 118, Route de Narbonne, 31062 Toulouse, Cedex 04, France
 bbourhim@cict.fr